

# Diagnostic Imaging Pathways - Scaphoid Fracture

## Population Covered By The Guidance

This pathway provides guidance on the imaging of adult patients with a suspected scaphoid fracture.

**Date reviewed: August 2013**

**Date of next review: August 2015**

**Published: August 2013**

## Quick User Guide


Move the mouse cursor over the **PINK** text boxes inside the flow chart to bring up a pop up box with salient points.

Clicking on the **PINK** text box will bring up the full text.

The relative radiation level (RRL) of each imaging investigation is displayed in the pop up box.

SYMBOL	RRL	EFFECTIVE DOSE RANGE
	None	0
	Minimal	< 1 millisieverts
	Low	1-5 mSv
	Medium	5-10 mSv
	High	>10 mSv

## Pathway Diagram



## Image Gallery

*Note: These images open in a new page*

**1 Scaphoid Fracture**



Image 1 (Plain Radiograph): Fracture of the scaphoid bone.

**2 Scaphoid Fracture**




Image 2 (Plain Radiograph): Undisplaced fracture of the scaphoid bone (arrows).

**3 Scaphoid Fracture**



Image 3 (Plain Radiograph): Non-union of scaphoid fracture. Note sclerosis at fracture line.

**4 Scaphoid Fracture**



Image 4 (Computed Tomography): Coronal image of the wrist showing fracture through the proximal waist of scaphoid (arrows).

## Teaching Points

- Early identification of the presence of a fracture and degree of displacement is important to prevent adverse complications such as avascular necrosis, non-union and osteoarthritis [1](#)
- The initial diagnostic modality for suspected scaphoid fracture is plain radiography, but up to 30% of fractures are radiographically occult at the time of presentation
- Based on current evidence, early MRI is most accurate in detecting occult scaphoid fracture, and has the advantage of simultaneously evaluating bone marrow abnormalities and surrounding soft tissue injuries
- Early imaging is preferred as it results in faster identification of fractures and other injuries and reduces unnecessary immobilisation. If early MRI is unavailable or contraindicated, CT is an alternative
- Depending on local resources, presumptive casting and repeat plain radiography in 7-10 days remains a specific alternative, but if clinical suspicion continues after negative follow-up radiographs, further imaging with MRI is required to exclude a scaphoid fracture
- Imaging modality choice will depend on patient age and preference, hand dominance, activity level and availability of MRI
- Evaluation of the diagnostic performance of different imaging modalities is limited by the lack of a consensus reference standard

## Plain Radiography

- Screening method of choice for evaluating clinically suspected scaphoid fracture
- Standard four view series (posteroanterior (PA), lateral, oblique and ulnar-deviated PA with cephalad angulation) is recommended. [2](#) Up to 30% of scaphoid fractures are radiographically occult at the time of presentation [3](#)
- Plain radiographs are even less sensitive for detecting other carpal fractures in patients with suspected scaphoid fracture [3](#)
- Specific (95%) but relatively insensitive (45%) in diagnosing displacement compared to arthroscopic assessment as the gold standard. [4](#) Presumptive casting and follow-up radiographs, when resorption at the fracture line may make previously occult fractures visible, has an estimated pooled 91.1% sensitivity and 99.8% specificity in the diagnosis of suspected scaphoid fractures on metaanalysis. [5](#) Reported sensitivities vary considerably, possibly due to poor interobserver agreement of follow-up radiograph findings [5,6](#)
- While useful if positive, follow-up radiographs cannot exclude fracture and further evaluation with MRI is recommended in the case of high clinical suspicion [5,6](#)

## Magnetic Resonance Imaging

- MRI is considered the best advanced imaging modality for suspected radiographically occult scaphoid fractures [3,5,7](#)
- A recent metaanalysis estimated a pooled 97.7% sensitivity and 99.8% specificity and excellent reliability for the diagnosis of suspected scaphoid fractures, more sensitive than CT or follow-up radiography and more specific than bone scan [5](#)
- There is mounting evidence that early MRI is marginally or comparably cost effective compared to conventional management with immobilisation and reassessment and may reduce the potentially large societal cost of unnecessary immobilisation, but is influenced by local availability of scanning resources [7-12](#)
- Hospitals with the facilities to implement early MRI have found it to have a high clinical impact [3,13,14](#)
- Allows in one test without ionising radiation the accurate detection of [7,13-15](#)
  - Occult scaphoid and other carpal bone fractures
  - Accompanying soft tissue injuries, a minority of which have the potential for significant morbidity, such as triangular fibrocartilage injuries and scapholunate ligament injuries [15-17](#)
  - Bone marrow abnormalities, such as avascular necrosis, and bone marrow oedema or 'bruising', which was reported to have a 2% risk of occult fracture development in one prospective study [18,19](#)
- Preoperative gadolinium-enhanced MRI can be used to evaluate the vascularity of the proximal pole in scaphoid non-unions which can help inform surgical decision to use a vascularised rather than standard non-vascularised bone graft. [20,21](#) However, it is relatively insensitive compared to intra-operative assessment as the gold standard and can not reliably predict prognosis in these patients. [22](#) Contrast enhanced MRI is also useful in post-operative monitoring [23](#)
- Disadvantages compared to CT include
  - Inferior to MDCT in depicting cortical involvement in occult scaphoid fractures [24](#)
  - Longer scanning time needed; is difficult for some patients where claustrophobia or pain may lead to increased risk of movement artefact

## Computed Tomography

- A recent metaanalysis estimated a pooled 85.2% sensitivity and 99.5% specificity for the diagnosis of suspected scaphoid fractures, inferior to MRI and associated with ionising radiation [5](#)
- May be used for suspected radiographically occult scaphoid fracture as an alternative when MRI is contraindicated or unavailable [3](#)
- Useful in
  1. Depicting occult scaphoid and other fractures of the wrist, where it is superior to plain radiographs, especially for capitate fractures, [3](#) and depicting cortical involvement in occult scaphoid fractures, where it is superior to MRI [24](#)
  2. Detecting scaphoid fracture displacement, [25](#) although compared to arthroscopic examination they remain only 77% sensitive. [4](#) This is important given it affects the decision to proceed to surgery over conservative management, and the risk of non-union [1](#)
  3. Surgical planning of complex fractures
  4. Diagnosis of union; less accurate at excluding non-union [26](#)
- CT arthrography is equal to MRI in detecting palmar and central segment tears and superior in detecting dorsal segment tears of the scapholunate and lunotriquetral ligaments in cadavers, [27](#) although it is limited by invasiveness

## Bone Scan

- A recent metaanalysis estimated a pooled 97.8% sensitivity and 93.5% specificity for the diagnosis of suspected scaphoid fracture [5](#)
- Similarly sensitive but less specific than MRI, and more sensitive but less specific than CT, in the detection of scaphoid fractures. [5](#) It is used for suspected radiographically occult scaphoid fracture as an alternative when MRI is contraindicated or unavailable
- A normal bone scan practically excludes a fractured scaphoid but specificity is limited by indiscriminate <sup>99m</sup>Tc-methylene disphosphonate uptake between underlying causes such as occult fracture, bone bruising, soft tissue injury, osteomyelitis or osteoarthritis [28](#)

## Ultrasonography

- Ultrasound is rapid and inexpensive but is user dependent, has variable sensitivity and specificity and relatively poor performance of both false positives and negatives. [7,29](#) It cannot exclude fracture, causes pain and it is difficult to image the proximal and distal poles of the scaphoid. Further delineation with other imaging is required
- Some studies have investigated the use of ultrasonography to triage patients to CT or MRI with clinical suspicion of radiographically occult scaphoid fractures [30,31](#)

## References

Date of literature search: May 2013

The search methodology is available on request. [Email](#)

References are graded from Level I to V according to the Oxford Centre for Evidence-Based Medicine, Levels of Evidence. [Download the document](#)

1. Singh HP, Taub N, Dias JJ. **Management of displaced fractures of the waist of the scaphoid:**

- meta-analyses of comparative studies.** Injury. 2012;43(6):933-9. (Level I evidence)
2. Expert Panel on Musculoskeletal Imaging; Rubin DA, Daffner RH, Weissman BN, Bennett DL, Blebea JS, et al. **American College of Radiology appropriateness criteria: acute hand and wrist trauma** [online publication]. Reston, VA; 2008 [cited 2013 April 4]. (Evidence based guidelines). [View the reference](#)
  3. Jorgsholm P, Thomsen NO, Besjakov J, Abrahamsson SO, Bjorkman A. **The benefit of magnetic resonance imaging for patients with posttraumatic radial wrist tenderness.** J Hand Surg Am. 2013;38(1):29-33. (Level II evidence)
  4. Buijze GA, Jorgsholm P, Thomsen NO, Bjorkman A, Besjakov J, Ring D. **Diagnostic performance of radiographs and computed tomography for displacement and instability of acute scaphoid waist fractures.** J Bone Joint Surg Am. 2012;94(21):1967-74. (Level II evidence)
  5. Yin ZG, Zhang JB, Kan SL, Wang XG. **Diagnostic accuracy of imaging modalities for suspected scaphoid fractures: meta-analysis combined with latent class analysis.** J Bone Joint Surg Br. 2012;94(8):1077-85. (Level I evidence)
  6. Low G, Raby N. **Can follow-up radiography for acute scaphoid fracture still be considered a valid investigation?** Clin Radiol. 2005;60(10):1106-10. (Level II evidence)
  7. Jenkins PJ, Slade K, Huntley JS, Robinson CM. **A comparative analysis of the accuracy, diagnostic uncertainty and cost of imaging modalities in suspected scaphoid fractures.** Injury. 2008;39(7):768-74. (Level II evidence)
  8. Patel NK, Davies N, Mirza Z, Watson M. **Cost and clinical effectiveness of MRI in occult scaphoid fractures: a randomised controlled trial.** Emerg Med J. 2013;30(3):202-7. (Level III evidence)
  9. Brooks S, Cicuttini FM, Lim S, Taylor D, Stuckey SL, Wluka AE. **Cost effectiveness of adding magnetic resonance imaging to the usual management of suspected scaphoid fractures.** Br J Sports Med. 2005;39(2):75-9. (Level II/III evidence)
  10. Dorsay TA, Major NM, Helms CA. **Cost-effectiveness of immediate MR imaging versus traditional follow-up for revealing radiographically occult scaphoid fractures.** AJR Am J Roentgenol. 2001;177(6):1257-63. (Level III evidence)
  11. Hansen TB, Petersen RB, Barckman J, Uhre P, Larsen K. **Cost-effectiveness of MRI in managing suspected scaphoid fractures.** J Hand Surg Eur Vol. 2009;34(5):627-30. (Level III evidence)
  12. Gooding A, Coates M, Rothwell A. **Cost analysis of traditional follow-up protocol versus MRI for radiographically occult scaphoid fractures: a pilot study for the Accident Compensation Corporation.** N Z Med J. 2004;117(1201):U1049. (Level III evidence)
  13. Khalid M, Jummani ZR, Kanagaraj K, Hussain A, Robinson D, Walker R. **Role of MRI in the diagnosis of clinically suspected scaphoid fracture: analysis of 611 consecutive cases and literature review.** Emerg Med J. 2010;27(4):266-9. (Level II evidence)
  14. Brydie A, Raby N. **Early MRI in the management of clinical scaphoid fracture.** Br J Radiol. 2003;76(905):296-300. (Level II evidence)
  15. McCullough NP, Smith FW, Cooper JG. **Early MRI in the management of the clinical scaphoid fracture.** Eur J Emerg Med. 2011;18(3):133-6. (Level II/III evidence)
  16. Magee T. **Comparison of 3-T MRI and arthroscopy of intrinsic wrist ligament and TFCC tears.** AJR Am J Roentgenol. 2009;192(1):80-5. (Level II evidence)
  17. Jorgsholm P, Thomsen NOB, Bjorkman A, Besjakov J, Abrahamsson S-O. **The incidence of intrinsic and extrinsic ligament injuries in scaphoid waist fractures.** J Hand Surg. 2010;35(3):368-74. (Level II/III evidence)
  18. Thavarajah D, Syed T, Shah Y, Wetherill M. **Does scaphoid bone bruising lead to occult fracture? A prospective study of 50 patients.** Injury. 2011;42(11):1303-6. (Level II evidence)
  19. La Hei N, McFadyen I, Brock M, Field J. **Scaphoid bone bruising--probably not the precursor of asymptomatic non-union of the scaphoid.** J Hand Surg Eur Vol. 2007;32(3):337-40. (Level II evidence)



20. Cerezal L, Abascal F, Canga A, Garcia-Valtuille R, Bustamante M, del Pinal F. **Usefulness of gadolinium-enhanced MR imaging in the evaluation of the vascularity of scaphoid nonunions.** AJR Am J Roentgenol. 2000;174(1):141-9. (Level II/III evidence)
21. Schmitt R, Christopoulos G, Wagner M, Krimmer H, Fodor S, van Schoonhoven J, et al. **Avascular necrosis (AVN) of the proximal fragment in scaphoid nonunion: is intravenous contrast agent necessary in MRI?** Eur J Radiol. 2011;77(2):222-7. (Level II evidence)
22. Megerle K, Worg H, Christopoulos G, Schmitt R, Krimmer H. **Gadolinium-enhanced preoperative MRI scans as a prognostic parameter in scaphoid nonunion.** J Hand Surg Eur Vol. 2011;36(1):23-8. (Level III evidence)
23. Dailiana ZH, Zachos V, Varitimidis S, Papanagiotou P, Karantanas A, Malizos KN. **Scaphoid nonunions treated with vascularised bone grafts: MRI assessment.** Eur J Radiol. 2004;50(3):217-24. (Level III evidence)
24. Memarsadeghi M, Breitenseher MJ, Schaefer-Prokop C, Weber M, Aldrian S, Gabler C, et al. **Occult scaphoid fractures: comparison of multidetector CT and MR imaging--initial experience.[Erratum appears in Radiology 2007;242(3):950].** Radiology. 2006;240(1):169-76. (Level II evidence)
25. Lozano-Calderon S, Blazar P, Zurakowski D, Lee SG, Ring D. **Diagnosis of scaphoid fracture displacement with radiography and computed tomography.** J Bone Joint Surg Am. 2006;88(12):2695-703. (Level III evidence)
26. Buijze GA, Wijffels MM, Guitton TG, Grewal R, van Dijk CN, Ring D. **Interobserver reliability of computed tomography to diagnose scaphoid waist fracture union.** J Hand Surg Am. 2012;37(2):250-4. (Level III evidence)
27. Schmid MR, Schertler T, Pfirrmann CW, Saupe N, Manestar M, Wildermuth S, et al. **Interosseous ligament tears of the wrist: comparison of multi-detector row CT arthrography and MR imaging.** Radiology. 2005;237(3):1008-13. (Level II evidence)
28. Fowler C, Sullivan B, Williams LA, McCarthy G, Savage R, Palmer A. **A comparison of bone scintigraphy and MRI in the early diagnosis of the occult scaphoid waist fracture.** Skeletal Radiol. 1998;27(12):683-7. (Level III evidence)
29. Yin Z-G, Zhang J-B, Kan S-L, Wang X-G. **Diagnosing suspected scaphoid fractures: a systematic review and meta-analysis.** Clin Orthop Relat Res. 2010;468(3):723-34. (Level I evidence)
30. Yildirim A, Unluer EE, Vandenberg N, Karagoz A. **The role of bedside ultrasonography for occult scaphoid fractures in the emergency department.** Ulus Travma Acil Cerrahi Derg. 2013;19(3):241-5. (Level III evidence)
31. Platon A, Poletti P-A, Van Aaken J, Fusetti C, Della Santa D, Beaulieu J-Y, et al. **Occult fractures of the scaphoid: the role of ultrasonography in the emergency department.** Skeletal Radiol. 2011;40(7):869-75. (Level II evidence)

## Information for Consumers

Information from this website	Information from the Royal Australian and New Zealand College of Radiologists' website
<a href="#">Consent to Procedure or Treatment</a> <a href="#">Radiation Risks of X-rays and Scans</a>	<a href="#">Computed Tomography (CT)</a> <a href="#">Gadolinium Contrast Medium</a>

<a href="#">Scaphoid Fracture (Suspected)</a>	<a href="#">Magnetic Resonance Imaging (MRI)</a>
<a href="#">Computed Tomography (CT)</a>	<a href="#">Plain Radiography/X-rays</a>
<a href="#">Magnetic Resonance Imaging (MRI)</a>	<a href="#">Radiation Risk of Medical Imaging During Pregnancy</a>
<a href="#">Plain Radiography (X-ray)</a>	<a href="#">Radiation Risk of Medical Imaging for Adults and Children</a>

## Copyright

© Copyright 2015, Department of Health Western Australia. All Rights Reserved. This web site and its content has been prepared by The Department of Health, Western Australia. The information contained on this web site is protected by copyright.

## Legal Notice

Please remember that this leaflet is intended as general information only. It is not definitive and The Department of Health, Western Australia can not accept any legal liability arising from its use. The information is kept as up to date and accurate as possible, but please be warned that it is always subject to change

## File Formats

Some documents for download on this website are in a Portable Document Format (PDF). To read these files you might need to download Adobe Acrobat Reader.

